

REFERENCES

- 1 Wade DT, Wood VA, Langton-Hewer R. Recovery after stroke—the first 3 months. *J Neurol Neurosurg Psychiatry* 1985;48:7-13.
- 2 Brocklehurst J, Andrews K, Richards B, et al. How much physical therapy for patients with stroke? *BMJ* 1978;1:45-9.
- 3 Kwakkel G, Wagenaar RC, Koelman TW, et al. Effects of intensity of rehabilitation after stroke. *Stroke* 1997;28:1550-6.
- 4 Langhorne P, Wagenaar RC, Partridge C. Physiotherapy after stroke: more is better? *Physiother Res Int* 1996;1:75-88.
- 5 Sunderland A, Tinson D, Bradley L, et al. Arm function after stroke. An evaluation of grip strength as a measure of recovery and a prognostic indicator. *J Neurol Neurosurg Psychiatry* 1989;52:1267-72.
- 6 Visintin M, Barbeau H, Lorner-Bitensky N, et al. A new approach to retrain gait in stroke patients through body weight support and treadmill stimulation. *Stroke* 1998;29:1122-8.
- 7 Van der Lee JH, Wagenaar RC, Lankhorst GJ, et al. Forced use of the upper extremity in chronic stroke patients: results from a single blind randomised controlled trial. *Stroke* 1999;30(11):2369-75.
- 8 Dean CM, Shepherd RB. Task-related training improves performance of seated reaching tasks after stroke. *Stroke* 1997;28(4):722-8.
- 9 Dickstein R, Hockerman S, Pillar T, et al. Stroke rehabilitation: three exercise therapy approaches. *Phys Ther* 1986;66:1233-8.
- 10 Loggigian MK, Samuels MS, Falconer J, et al. Clinical exercise trial for stroke patients. *Arch Phys Med Rehabil* 1983;64:364-7.
- 11 Lord JP, Hall K. Neuromuscular reeducation versus traditional programs for stroke rehabilitation. *Arch Phys Med Rehabil* 1986;67:88-91.
- 12 Ernst E. A review of stroke rehabilitation and physiotherapy. *Stroke* 1990;21:1081-5.
- 13 Ashburn A. Physiotherapy in the rehabilitation of stroke: a review. *Clin Rehabil* 1993;7:337-45.
- 14 Wagenaar RC, Meijer OG, van Wieringen CW, et al. The functional recovery of stroke: a comparison between neuro-developmental treatment and the Brunstrom method. *Scand J Rehabil Med* 1990;22:1-8.
- 15 Nilsson L, Nordholm L. Physical therapy in stroke rehabilitation: bases for Swedish physiotherapist's choice of treatment. *Physiother Theory Pract* 1992;8:49-55.
- 16 Davidson I, Waters K. Physiotherapists working with stroke patients. *Physiotherapy* 2000;86(2):69-80.
- 17 Bobath B. *Adult hemiplegia. Evaluation and treatment*, 3rd ed. Oxford: Heinemann Medical, 1990.
- 18 Davies P. *Right in the middle*. Heidelberg: Springer, 1991.
- 19 Carr JH, Shepherd RB. *A motor relearning programme for stroke*. London: Heinemann Physiotherapy, 1987.
- 20 Carr JH, Shepherd RB. *Neurological rehabilitation: optimising motor performance*. Oxford: Butterworth-Heinemann, 1998.
- 21 van Vliet PM, Lincoln NB, Robinson E. Comparison of the content of two physiotherapy approaches for stroke. *Clin Rehabil* 2000;15:398-414.
- 22 Langhammer B, Stanghelle JK. Bobath or Motor Relearning Programme? A comparison of two different approaches of physiotherapy in stroke rehabilitation: a randomized controlled study. *Clin Rehabil* 2000;14:361-9.

- 23 Paci M. Physiotherapy based on the Bobath concept for adults with post-stroke hemiplegia: a review of effectiveness studies. *J Rehabil Med* 2003;35:2-7.
- 24 Bamford J, Sandercock P, Dennis M, et al. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet* 1991;337:1521-6.
- 25 Davies PM. Steps to follow: a guide to the treatment of adult hemiplegia. Heidelberg: Springer, 1985.
- 26 Carr JH, Shepherd RB. Movement Science: foundations for physical therapy in rehabilitation. London: Heinemann Physiotherapy, 1987.
- 27 Carr JH, Shepherd RB. A motor learning model for stroke rehabilitation. *Physiotherapy* 1989;89:372-80.
- 28 Carr JH, Shepherd RB. Reflections on physiotherapy and the emerging science of movement rehabilitation. *Aust J Physiother*, 1994;40th Jubilee, 39-47.
- 29 Lincoln NB, Leadbitter D. Assessment of motor function in stroke patients. *Physiotherapy* 1979;65:48-51.
- 30 Carr JH, Shepherd RB, Nordholm L, et al. Investigation of a new motor assessment scale for stroke patients. *Phys Ther* 1985;65(2):175-80.
- 31 Turton AJ, Fraser CM. A test battery to measure the recovery of voluntary movement following stroke. *Int Rehabil Med* 1986;8:74-9.
- 32 Wade DT, Hower RL, Skilbeck CE, et al. Stroke: a critical approach to diagnosis, treatment and management. London: Chapman and Hall, 1985.
- 33 Bohannon RW, Smith MB. Interrater reliability of a Modified Ashworth Scale of muscle spasticity. *Phys Ther* 1987;2:206-7.
- 34 Jackson J, Crow L. The reliability of sensory assessments in hemiplegia. Paper presented at the Eleventh Congress of the World Confederation for Physical Therapy, 1991, London.
- 35 Collin C, Wade DT, Davies S, et al. The Barthel ADL Index, a reliability study. *Int Disabil Stud* 1988;10:61-3.
- 36 Nouri FM, Lincoln NB. An extended activities of daily living scale for stroke patients. *Clin Rehabil* 1987;1:301-5.
- 37 Syder D, Body R, Parker M, et al. Sheffield Screening Test for Acquired Language Disorders. Windsor: NFER Nelson, 1993.
- 38 Rey A. Le test, de copie de figure complexe. Paris: Editions Centre de Psychologie Applique, 1959.
- 39 Wilson B, Cockburn J, Halligan PW. Behavioural Inattention Test. Titchfield, Hampshire: Thames Valley Test Company, 1987.
- 40 Wilson BA, Cockburn J, Baddeley A. The Rivermead Behavioural Memory Test, Bury St. Edmunds, Suffolk: Thames Valley Test Company, 1985.
- 41 Juby LC, Lincoln NB, Berman P. The effect of a stroke rehabilitation unit on functional and psychological outcome: a randomised controlled trial. *Cerebrovasc Dis* 1996;6:106-10.
- 42 Matthews JNS, Altman DG, Campbell MJ, et al. Analysis of serial measurements in medical research. *BMJ* 1990;300:230-5.
- 43 Platz T, Winter T, Muller N, et al. Arm ability training for stroke and traumatic brain injury patients with mild arm paresis: a single-blind, randomized, controlled trial. *Arch Phys Med Rehabil* 2001;82:961-8.
- 44 Feys HM, Weerdt WD, Selz BE, et al. Effect of a therapeutic intervention for the hemiplegic upper limb in the acute phase after stroke. A single-blind, randomized, controlled multicenter trial. *Stroke* 1998;29:785-92.
- 45 Sunderland A, Tinson D, Bradley L, et al. Enhanced physical therapy for arm function after stroke. *J Neurol Neurosurg Psychiatry* 1992;55:530-5.

46 Kwakkel G, Wagenaar RC, Twisk JWR, et al. Intensity of leg and arm training after primary middle-cerebral artery stroke: a randomised trial. *Lancet* 1999;354:191-6.

47 Ada L, Canning C. Key issues in neurological physiotherapy. London: Heinemann Medical, 1990.

48 Carr JH, Shepherd RB. Spasticity: research findings and implications for intervention. *Physiotherapy* 1995;81:421-9.

49 Kidd G, Lawes N, Musa I. Understanding neuro-muscular plasticity. London: Edward Arnold, 1992.

50 Mulder T. A process-oriented model of human motor behaviour: toward a theory based rehabilitation approach. *Phys Ther* 1991;71:157-64.

51 Rottke K. From reflex to skill—the training of co-ordination. *Arch Phys Med Rehabil* 1980;61:551-61.

52 Lincoln NB, Parry RH, Vass CD. Randomized, controlled trial to evaluate increased intensity of physiotherapy treatment of arm function after stroke. *Stroke* 1999;30:573-9.

508 van Vliet, Lincoln, Foxall

www.jnnp.com

Downloaded from jnnp.bmj.com on 7 December 2006

- ◆ P.Davies "passo dopo passo, emiplegia, ricominciare" ed.Springer; Schmidt "
- ◆ apprendimento motorio e prestazione" ed.società sportiva Roma;
- ◆ Kandel "principi di neuroscienze" ed. ambrosiana Milano.